



Aviax[®] Plus

DOSAGE FORM:

Premix

COMPOSITION:

Semduramicin 30 g/kg, Nicarbazin 80 g/kg. The two active ingredients in Aviax[®] Plus are granulated together to ensure optimum efficacy.

INDICATIONS:

For the prevention of coccidiosis in broiler chickens caused by *Eimeria tenella*, *E. acervulina*, *E. maxima*, *E. brunetti*, *E. necatrix* and *E. mivati/mitis*.

WARNINGS:

- Do not feed to laying hens.
- Withdrawal period for meat: One day before slaughter.
- Keep out of reach of children, uninformed persons and animals.
- Although this remedy has been extensively tested under a large variety of conditions, failure thereof may ensue as a result of a wide range of reasons. If this is suspected, seek veterinary advice and notify the registration holder.

DOSAGE AND DIRECTIONS FOR USE:

Thoroughly mix 500 g of Aviax[®] Plus into a metric ton of feed. If high coccidiosis challenge levels are present, the inclusion level of Aviax[®] Plus can be increased to 550 or 600 g per ton of feed.

PRESENTATION:

25 kg bags.

STORAGE INSTRUCTIONS:

Store in a cool, dry place below 25 °C

REGISTRATION NUMBER:

G3693 (Act 36/1947)

REGISTRATION HOLDER:

Phibro Animal Health (Pty) Ltd.
Company Reg. No.: 2000/004664/07.
P O Box 5388, Rietvalleirand, Gauteng 0174
63 Regency Drive, Block B Technolink,
Route 21 Corporate Park, Irene.
T: +27 12 4500 603 • F: +27 12 345 5357
National Call Centre: 086 11 11 900
www.phibro.co.za
www.pahc.com
www.phibro-vaccines.com