

Coryza AKKO + 221

DESCRIPTION:

An inactivated, water-in-oil emulsion, vaccine for the immunization of layers and breeder fowls against Infectious Coryza.

COMPOSITION:

Each dose of the vaccine contains inactivated *Avibacterium paragallinarum* (previously *Haemophilus paragallinarum*) AKKO and 221 strains. Minimum titre per dose: $10^{8.6}$ CFU / dose of serotype C, strain AKKO, and $10^{8.4}$ CFU / dose of serotype A, strain 221.

DIRECTIONS FOR USE: Use only as directed.

- Allow the vaccine to gradually reach room temperature before commencing with vaccination.
- The emulsion should be shaken well before, and at regular intervals during, use.
- **The vaccine is to be administered to fowls by subcutaneous injection only.**
- It is advisable that the initial vaccination should be at 7 - 8 weeks of age, and a second booster injection 3 - 4 weeks after primary vaccination and before point of lay.

Consult a veterinarian on the best vaccination programme schedule for your circumstances.

RECOMMENDED DOSAGE:

Inject 0,5 ml per fowl subcutaneously on the back between the wings lateral to the spine.

PRECAUTIONS:

See product label.

WARNINGS:

See product label for full details.

STORAGE:

Store between 2 °C and 8 °C. Protect from light.

Do not freeze.

PRESENTATION:

250 ml or 500 ml containing 500 or 1 000 doses respectively.

REGISTRATION NUMBER:

G3800 (Act 36/1947)

REGISTRATION HOLDER:

Phibro Animal Health (Pty) Ltd.

Company Reg. No.: 2000/004664/07.

P O Box 5388, Rietvalleirand, Gauteng 0174

63 Regency Drive, Block B Technolink,

Route 21 Corporate Park, Irene.

T: +27 12 4500 603 • F: +27 12 345 5357

National Call Centre: 086 11 11 900

www.phibro.co.za

www.pahc.com

www.phibro-vaccines.com