

CoxAbic®

DESCRIPTION:

A Coccidiosis vaccine.

COMPOSITION:

The vaccine contains a purified antigen isolated from the gametocyte stage of *Eimeria maxima* in an oil emulsion.

INDICATIONS:

An water-in-oil emulsion vaccine for vaccination of breeder pullets to protect their offspring against coccidiosis.

DIRECTIONS FOR USE: Use only as directed.

The vaccine is administered to breeding pullets by intramuscular injection in the breast muscle. For best results, two vaccinations should be given: The first at 12 - 16 weeks of age and the second just prior to point of lay (approximately between 18 and 20 weeks) at an interval of at least 3 to 4 weeks. CoxAbic® can be given at the same time as other vaccines.

The progeny of the vaccinated flock should be protected throughout the whole laying period, provided that the antibody titres in the breeder hens are found to be above S/P of 0,4 in the ELISA test.

The vaccination should be carried out at a time when the birds are kept as quiet as possible (i.e. limited lighting).

It is advisable to consult your veterinarian for the suitable vaccination programme for your birds.

The vaccine should be allowed to gradually reach room temperature (20 - 25 °C) before use.

Shake the bottle well before use and at regular intervals during use.

IMPORTANT POINTS CONCERNING THE BROILERS HATCHED FROM VACCINATED HENS

- Ensure that the broilers only receive feed containing no coccidiostats. This is especially critical in the first three weeks of life.
- Samples of all feed deliveries should be taken and kept for reference purposes until the flock has been slaughtered. Should it be found that coccidiostats have been included, the flock should be raised further on coccidiostats.
- Most approved growth promoters can be safely used in broiler feed.

- Potentially coccidiocidal drugs should not be administered to broilers as they may have a negative effect on the development of immunity.

RECOMMENDED DOSAGE:

Inject all the fowls in the flock with 0,5 ml intramuscularly (into the breast muscle).

The needle should be pointed in the direction of the fowl's head to prevent the needle from entering the body cavity.

PRECAUTIONS:

- Eggs from CoxAbic® vaccinated breeder hens should be separated from eggs laid by unvaccinated breeder hens and incubated separately.
- Chicks from vaccinated breeder hens should be fed with feed completely free of coccidiostats and coccidiocides such as sulphonamides, nitrofurans, etc.
- See label for full precautions.

WARNINGS:

See label for warnings.

STORAGE:

Keep refrigerated between 2 °C and 8 °C.

Do not freeze. Use within expiry date.

PRESENTATION:

500 ml (1 000 doses)

REGISTRATION NUMBER:

G3024 (Act 36/1947)

REGISTRATION HOLDER:

Phibro Animal Health (Pty) Ltd.

Company Reg. No.: 2000/004664/07.

P O Box 5388, Rietvalleirand, Gauteng 0174

63 Regency Drive, Block B Technolink,

Route 21 Corporate Park, Irene.

T: +27 12 4500 603 • F: +27 12 345 5357

National Call Centre: 086 11 11 900

www.phibro.co.za

www.pahc.com

www.phibro-vaccines.com